

Ionic criticality : an exactly soluble model

Jean-Noël Aqua and Michael E. Fisher

Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742, USA

(Dated: February 2, 2008)

Gas-liquid criticality in ionic fluids is studied in exactly soluble spherical models that use interlaced sublattices to represent hard-core *multicomponent* systems. Short range attractions in the uncharged fluid drive criticality but charged ions do not alter the universality class. Debye screening remains exponential *at* criticality in charge-symmetric 1:1 models. However, *asymmetry* couples charge and density fluctuations in a direct manner: the charge correlation length then diverges precisely as the density correlation length and the Stillinger-Lovett rule is violated *at* criticality.

PACS numbers: 64.60.Fr, 61.20.Qg, 05.50.+q, 64.70.Fx

The nature of gas-liquid (or, more generally fluid-fluid) criticality in systems in which long-range ionic interactions play a significant role has been a focus of attention since still-puzzling experiments questioned the appropriate universality class [1]. Beyond further experiments [1], numerous theoretical [2, 3, 4, 5] and computational [6] studies have been reported; however, basic questions remain open. Certainly, the character of criticality depends on the range of the interactions: One expects an Ising critical point in a fluid with short-range couplings but mean-field behavior for interactions of sufficiently long-range. So, might the introduction of ions interacting via long-range Coulomb forces destroy an Ising-type critical point? Coulomb interactions are exponentially screened in a conducting classical fluid, as proved rigorously at low densities [7]. Charge fluctuations in a fluid of $\mathcal{S}$ species of charges q_σ and valences $z_\sigma = q_\sigma/q$ (where q is an elementary charge), thus decay over a few Debye screening lengths ξ_D , where

$$\xi_D^{-2}(T, \rho) = 4\pi\rho q^2 \sum_{\sigma=1}^{\mathcal{S}} z_\sigma^2 x_\sigma / k_B T, \quad (1)$$

in which $\rho = \sum_\sigma \rho_\sigma$ is the overall density while the mole fractions are $x_\sigma = \rho_\sigma/\rho$. But, does exponential screening on this scale hold near and at criticality?

Indeed, a major open issue is the behavior of the pairwise *charge* correlations near the critical point, where the *density* fluctuations diverge strongly. With

$$G_{\sigma\tau}(\mathbf{r}) = \langle \rho_\sigma(\mathbf{0}) \rho_\tau(\mathbf{r}) \rangle - \rho_\sigma \rho_\tau, \quad (2)$$

the correlation functions, G_{NN} , G_{ZZ} , and G_{NZ} for the density, charge, and charge-density, are [4]

$$G_{XY}(\mathbf{r}; T, \rho) = \sum_{\sigma, \tau} q_\sigma^{\vartheta_X} q_\tau^{\vartheta_Y} G_{\sigma\tau}(\mathbf{r}; T, \rho), \quad (3)$$

where X and Y may be N or Z while $\vartheta_N = 0$ and $\vartheta_Z = 1$. Except at the critical point (T_c, ρ_c) , itself, we may suppose that the corresponding structure factor, S_{NN} , has the small $k = |\mathbf{k}|$ expansion

$$S_{NN}(\mathbf{k})/S_{NN}(\mathbf{0}) = 1 + \sum_{p \geq 1} (-)^p \xi_{N,p}^{2p}(T, \rho) k^{2p}. \quad (4)$$

Near criticality, $S_{NN}(\mathbf{0}; T, \rho_c)$ diverges as $1/t^\gamma$ when $t \equiv (T - T_c)/T_c \rightarrow 0+$, while the length $\xi_{N,\infty}$ characterizing the exponential decay of $G_{NN}(\mathbf{r}; T, \rho_c)$ diverges as ξ_N^0/t^ν (where short-range forces have been assumed). At criticality, density fluctuations are long-ranged and $S_{NN}(\mathbf{k}; T_c, \rho_c) \sim 1/k^{2-\eta}$.

By contrast, the charge structure factor should obey

$$S_{ZZ}(\mathbf{k}) = 0 + \xi_{Z,1}^2 k^2 - \sum_{p \geq 2} (-)^p \xi_{Z,p}^{2p}(T, \rho) k^{2p}, \quad (5)$$

where the first vanishing term results from electroneutrality reflecting the *internal* screening in an ionic fluid, while the Stillinger-Lovett sum-rule [8, 9]

$$\xi_{Z,1}(T, \rho) = \xi_D(T, \rho), \quad (6)$$

characterizes the screening of *external* charges. Does this hold near and at criticality? Finally, we focus also on the charge correlation length $\xi_{Z,\infty}(T, \rho)$, that specifies the exponential decay of $G_{ZZ}(\mathbf{r})$ when $r \rightarrow \infty$. How does $\xi_{Z,\infty}$ vary when $\xi_{N,\infty}$ diverges near criticality?

To obtain insight into these questions, we study *multicomponent* lattice gas generalizations of the spherical model [10, 11, 12] specifically designed to represent hard-core interactions and thus avoid the mutual “annihilation” of oppositely charged ions on the same site. This crucial feature, which (in contrast to [11]) allows gas-liquid criticality to survive in the presence of Coulomb interactions, is accomplished by using a set of equivalent interlacing sublattices (with sites i at $\mathbf{R}_i^\sigma$ with spacing a), one for each of the $\sigma = 1, 2, \dots, \mathcal{S}$ distinct particle species [13]. Thereby unlike charges cannot approach closer than an effective hard-core diameter a_0 : see, e.g., Fig. 1.

To specify the models more fully, consider a multicomponent grand canonical lattice gas with site occupancy variables $n_i^\sigma = 0, 1$; this is equivalent to an Ising magnet with spins $s_i^\sigma = 2n_i^\sigma - 1 = \pm 1$, subject to fields h_σ (linearly related to the chemical potentials, μ_σ). For attractive interparticle potentials, the corresponding spin-spin couplings, $J_{\sigma\tau}(\mathbf{R})$, at lattice separations $\mathbf{R}$, are positive. We decompose these couplings as

$$J_{\sigma\tau}(\mathbf{R}) = J_{\sigma\tau}^0(\mathbf{R}) - \frac{1}{4} q_\sigma q_\tau \varphi^C(\mathbf{R}), \quad (7)$$




FIG. 1: Interlaced + and - sc sublattices with $a_0 = \frac{1}{2}\sqrt{3}a$.

where $J_{\sigma\tau}^0(\mathbf{R})$ represents short-range or, more generally, integrable interparticle interactions which we suppose suffice to drive gas-liquid criticality (even if the charges q_σ vanish). We take the d -dimensional Coulomb potential, $\varphi^C(\mathbf{R}) \sim 1/R^{d-2}$ ($d > 2$), as the solution of an appropriate discrete Laplace equation [14] (with, for convenience, a uniform neutralizing background so that electroneutrality, $\sum_\sigma x_\sigma q_\sigma = 0$, must be imposed). Fourier transforms, $\hat{J}_{\sigma\tau}(\mathbf{k})$, are defined by summing over one sub-lattice, with Brillouin zone $\mathcal{B}$, and it is useful to introduce

$$\Delta J_{\sigma\tau}(\mathbf{k}) \equiv \frac{1}{2}[\hat{J}_{\sigma\tau}(\mathbf{0}) - \hat{J}_{\sigma\tau}(\mathbf{k})]. \quad (8)$$

To model a simple 1:1 electrolyte one needs only $\mathcal{S} = 2$ components, say + and - with $q_\pm = \pm q$. We may then identify basic energy and range scales, j_0 and R_0 , via

$$\hat{J}_{+-}^0(\mathbf{0}) \equiv k_B T_0 \equiv 2j_0 > 0, \quad (9)$$

and, assuming short-range isotropic nonionic couplings,

$$\Delta J_{++}^0 + 2\Delta J_{+-}^0 + \Delta J_{--}^0 \approx 2j_0 R_0^2 k^2 > 0, \quad (10)$$

as $k \rightarrow 0$. Now, if $J_{++}(\mathbf{R}) = J_{--}(\mathbf{R})$, the model is (fully) *charge symmetric* (as is the well known continuum Restricted Primitive Model or RPM [1, 2, 3, 6]). A suitable *charge asymmetry* parameter is then [14]

$$\delta_J = \max_{\mathbf{k} \in \mathcal{B}} |\Delta J_{++}(\mathbf{k}) - \Delta J_{--}(\mathbf{k})| / k_B T_0, \quad (11)$$

which might, e.g., be used to represent distinct ionic sizes, a_+ and a_- [6(a)]. As the simplest “*Basic Ionic Model*” it suffices to take only nearest neighbor couplings, $J_{+-}^0 > 0$ and $J_{++}^0 = -J_{--}^0$; then one has $j_0 = 2^{d-1}J_{+-}^0$ and $\delta_J = d|J_{++}^0|/2^{d-2}|J_{+-}^0|$. Finally, as a dimensionless measure of the relative strength of the Coulomb interactions, it is helpful to introduce the *ionicity* [15]

$$\mathcal{I}_0 = q^2 / a^{d-2} k_B T_0. \quad (12)$$

Of course, even this Basic Ionic Model is insoluble for $d \geq 2$. Accordingly, in the standard way [10, 11, 12],

we “sphericalize” these multicomponent models by taking the spins s_i^σ as unbounded continuous variables subject only to $\mathcal{S}$ *spherical constraints*, $\langle s_\sigma^2 \rangle = 1$, enforced with the aid of $\mathcal{S}$ Lagrange multipliers which, for later convenience, we write as $\lambda_\sigma + \frac{1}{2}\hat{J}_{\sigma\sigma}(\mathbf{0})$. In full generality, the singular part of the total free energy $f[T, \mathbf{h}; \boldsymbol{\lambda}(T, \mathbf{h})]$ in the thermodynamic limit, is then [14]

$$f_s(T; \boldsymbol{\lambda}) = \frac{1}{2} k_B T \int_{\mathbf{k}} \ln \{ |\boldsymbol{\Lambda}(\mathbf{k}; \boldsymbol{\lambda})| / (k_B T)^{\mathcal{S}} \}, \quad (13)$$

where $\int_{\mathbf{k}} \equiv (a/2\pi)^d \int_{\mathbf{k} \in \mathcal{B}} d^d \mathbf{k}$ and $\mathbf{h} = (h_\sigma)$ and $\boldsymbol{\lambda} = (\lambda_\sigma)$, while the $\mathcal{S} \times \mathcal{S}$ matrix $\boldsymbol{\Lambda}$ has elements

$$\Lambda_{\sigma\tau} = [\lambda_\sigma + \Delta J_{\sigma\sigma}(\mathbf{k})] \delta_{\sigma\tau} - \frac{1}{2}(1 - \delta_{\sigma\tau}) \hat{J}_{\sigma\tau}(\mathbf{k}). \quad (14)$$

The field-dependent contribution to the free energy is

$$f_h = -\frac{1}{4} \langle \mathbf{h} | \boldsymbol{\Lambda}^{-1}(\mathbf{0}; \boldsymbol{\lambda}) | \mathbf{h} \rangle \quad \text{with} \quad \mathbf{h} = 2\boldsymbol{\Lambda}(\mathbf{0}; \boldsymbol{\lambda}) \mathbf{m}, \quad (15)$$

where $\mathbf{m} = (m_\sigma)$ and $m_\sigma = \langle s_\sigma \rangle = 2\langle n_\sigma \rangle - 1$ so that $\rho_\sigma = \frac{1}{2} \rho_\sigma^{max} (1 + m_\sigma)$. Finally, the Lagrange multipliers $\boldsymbol{\lambda}(T, \mathbf{h})$ are determined implicitly via the $\mathcal{S}$ spherical constraints $\langle s_\sigma^2 \rangle = (\partial f / \partial \lambda_\sigma) = 1$. These results are valid while the eigenvalues of $\boldsymbol{\Lambda}$ remain positive; the vanishing of any one of them signals a thermodynamic singularity.

For brevity hereon we focus on the two-species, 1:1 case (with $q_\pm = \pm q$), the + and - ions residing on one of two hypercubic sublattices displaced by $\frac{1}{2}(a, a, \dots, a)$: see Fig.1 for $d = 3$. The eigenvalues of $\boldsymbol{\Lambda}$, to be called Λ_z and Λ_N for reasons soon to be evident, are then simply the + and - roots of the quadratic equation $|\boldsymbol{\Lambda} - x\mathbf{I}| = 0$,

$$\Lambda_x(\mathbf{k}; \boldsymbol{\lambda}) = \bar{\lambda} + \Delta \bar{J}(\mathbf{k}) - (-)^{\vartheta_x} \mathcal{D}(\mathbf{k}; \boldsymbol{\lambda}) \quad \text{with} \quad (16)$$

$$\mathcal{D}^2(\mathbf{k}; \boldsymbol{\lambda}) \equiv [\lambda^\dagger + \Delta J^\dagger(\mathbf{k})]^2 + \frac{1}{4} \hat{J}_{+-}^2(\mathbf{k}), \quad (17)$$

where, for each variable g_+ , g_- , g_{++} , etc., we have introduced the mean $\bar{g} = \frac{1}{2}(g_+ + g_-)$ and difference $g^\dagger = \frac{1}{2}(g_+ - g_-)$, while the square root is chosen so that $\mathcal{D}(\mathbf{0}; \boldsymbol{\lambda}) > 0$ and $\Lambda_x(\mathbf{k})$ is analytic.

A pivotal result now transpires [14], namely, the linear decomposition of the structure factors via

$$\frac{S_{xy}(\mathbf{k}; \boldsymbol{\lambda})}{k_B T / 4\rho a^d} = \frac{B_{xy}^N(\mathbf{k}; \boldsymbol{\lambda})}{\Lambda_N(\mathbf{k}; \boldsymbol{\lambda})} + \frac{B_{xy}^Z(\mathbf{k}; \boldsymbol{\lambda})}{\Lambda_Z(\mathbf{k}; \boldsymbol{\lambda})}, \quad (18)$$

in which $B_{NN}^N = B_{ZZ}^Z = 1 - B_{NN}^Z = 1 - B_{ZZ}^N$, while

$$B_{NN}^N(\mathbf{k}; \boldsymbol{\lambda}) = \frac{1}{2} + \frac{1}{4} |\hat{J}_{+-}(\mathbf{k})| / \mathcal{D}(\mathbf{k}; \boldsymbol{\lambda}), \quad (19)$$

and $B_{NZ}^N + B_{NZ}^Z = 0$. In the *charge symmetric* cases, all the $g^\dagger$ variables vanish by definition, so $B_{NN}^N \equiv 1$ and $B_{NN}^Z = B_{ZZ}^N \equiv 0$: this implies that S_{NN} is entirely governed by Λ_N , and, likewise, S_{ZZ} by Λ_Z , so justifying the notation. Conversely, in *nonsymmetric* models, both Λ_N and Λ_Z contribute to *all* the structure functions.

Finally, the spherical constraints reduce, first, to

$$k_B T \mathcal{J}_d(\boldsymbol{\lambda}) + \bar{h}^2 / 4\lambda^2 = 1 \quad \text{with} \quad \lambda = \bar{\lambda} - j'_0, \quad (20)$$

where $j'_0 = \frac{1}{2}\hat{J}_{+-}(\mathbf{0})$ [14], while the basic integral

$$\mathcal{J}_d(\boldsymbol{\lambda}) = \frac{1}{4} \int_{\mathbf{k}} [\Lambda_N^{-1}(\mathbf{k}; \boldsymbol{\lambda}) + \Lambda_Z^{-1}(\mathbf{k}; \boldsymbol{\lambda})], \quad (21)$$

becomes singular, typically as $\Delta\mathcal{J}_d \sim -\lambda^{1/\gamma}$, when λ ($\sim t^\gamma$) vanishes, where γ is the appropriate critical exponent [10, 12, 14]; lastly $\lambda^\dagger$ is given (implicitly) by

$$\lambda^\dagger \int_{\mathbf{k}} 1/|\Lambda(\mathbf{k}; \boldsymbol{\lambda})| = - \int_{\mathbf{k}} \Delta J^\dagger(\mathbf{k})/|\Lambda(\mathbf{k}; \boldsymbol{\lambda})|. \quad (22)$$

Recall, however, that $\lambda^\dagger$ vanishes identically in charge symmetric cases: more generally, this result relates $\lambda^\dagger$ to the asymmetry parameter δ_J [see Eq. (11)] via

$$\lambda_c^\dagger \approx c_\delta[J_{\sigma\tau}] \delta_J, \quad (23)$$

where, however, $c_\delta[J_{\sigma\tau}]$ might vanish “accidentally”.

We are now in a position to answer, with explanations, the questions posed after Eqns. (1)-(6). For specific numerical results we will invoke the *Basic Ionic Spherical Model* (BISM), i.e., the sphericalized version of the $d = 3$, $\mathcal{S} = 2$, 1:1 model with $\rho_+ = \rho_- = \frac{1}{2}\rho$ and nearest-neighbor interactions as set out with Eqns. (9)-(12). When $\delta_J = \mathcal{I}_0 = 0$ one readily finds from the vanishing of $\Lambda_N(\mathbf{0})$ when $\lambda \rightarrow 0$ [10, 11, 12, 14] that standard spherical-model criticality and scaling pertains, with exponents $\beta = \frac{1}{2}$ and $\gamma = 1 - \alpha = (2 - \eta)\nu$, where short-range (non-Coulomb) interactions [16] lead to $\eta = 0$ and $\gamma = \max\{2/(d-2); 1\}$ (for $d \geq 2$).

For $\mathcal{I}_0$ and δ_J not too large, the *same* situation prevails—contrary to speculations (for hard-core continuum ionic models) that lack of symmetry might lead to mean-field criticality [3]. This follows most directly from the small $\mathbf{k}$ behavior of the eigenvalues, namely,

$$\begin{aligned} \Lambda_N(\mathbf{k}; \boldsymbol{\lambda}) &= \lambda + j_0 R_N^2(\boldsymbol{\lambda}) k^2 + \mathcal{O}(k^4), \\ \Lambda_Z(\mathbf{k}; \boldsymbol{\lambda}) &= (S_d q^2 / 4a^d) [1/k^2 + R_Z^2 + \mathcal{O}(k^2)], \end{aligned} \quad (24)$$

where S_d is the area of a unit d -sphere, while [14]

$$\begin{aligned} R_N^2(\boldsymbol{\lambda}) &= R_0^2 - S'_d a^2 \mathcal{I}_0 - 2a^2 \lambda^{\dagger 2} / S_d \mathcal{I}_0 j_0^2, \\ R_Z^2(\boldsymbol{\lambda}, \hat{\mathbf{k}}) &= 2a^2 (\lambda + j'_0) / S_d \mathcal{I}_0 j_0 + a^2 \Sigma_4(\hat{\mathbf{k}}), \end{aligned} \quad (25)$$

in which $S'_3 = \pi/144$ and $\Sigma_4 \equiv \frac{1}{48} \sum_{\alpha} k_{\alpha}^4 / k^4$. The crucial feature, following directly from (16), is that the Coulomb singularity, characterized by $\hat{\varphi}^c(\mathbf{k}) \sim 1/k^2$, cancels out of Λ_N exactly thanks to electroneutrality. (The absence of this possibility for $\mathcal{S} = 1$ results in the destruction of gas-liquid criticality by Coulomb interactions [11(b)].)

To ensure the stability of the critical point, one also needs [14]: (i) $\mathcal{I}_0 < \mathcal{I}_d^{max}$, with $\mathcal{I}_3^{max} = \frac{96}{11}\pi \simeq 2.77$; (ii) $R_N^2 > 0$, which restricts $\mathcal{I}_0$ and δ_J to the interior of an ellipsoid with a vertex at $\delta_J = \mathcal{I}_0 = 0$, which, for $d = 3$, is $\frac{1}{72}\delta_J^2 + [\frac{1}{72}\pi\mathcal{I}_0 - (R_0^2/a^2)]^2 < R_0^4/a^4$; and (iv) the absence of competing minima in $\Lambda_N(\mathbf{k})$,

which is satisfied for sufficiently small δ_J , specifically by $\delta_J < 1 - \frac{1}{12}\pi\mathcal{I}_0$ in the BISM. The solution $\lambda^\dagger(\lambda)$ of (22) then varies smoothly when $\lambda \rightarrow 0$ and (23) applies. By (20) criticality is restricted to $\bar{h} = 0$ and occurs at $k_B T_c = 1/\mathcal{J}_d(0, \lambda_c^\dagger)$ and $\rho_c = a^{-d} = \frac{1}{2}\rho_{max}$. For the BISM, we find $T_c \approx T_0/K_{bcc}$, to lowest order in δ_J and $\mathcal{I}_0$, with $K_{bcc} \simeq 1.39$ [10]. It transpires [14] that $T_c(\delta_J)$ is a *decreasing* function of the asymmetry in accord with recent simulations of hard-core continuum electrolyte models [6(a)] that, however, contradict various approximate theories. Furthermore, a term varying as $\mathcal{I}_0^{3/2}$ (in $d = 3$) appears [14], in accord with [15] and in analogy with [17].

As regards the density correlation lengths, (18) yields

$$\xi_{N,1}(T, \rho) = R_N[\boldsymbol{\lambda}(T, \rho)] [j_0/\lambda(T, \rho)]^{1/2}, \quad (26)$$

for all acceptable $\mathcal{I}_0$ and δ_J ; when $\rho = \rho_c$, this diverges as ξ_N^0/t^ν . Furthermore, all higher moments [see (4)], including the “true” correlation length $\xi_{N,\infty}$ [18], satisfy $\xi_{N,p}/\xi_{N,1} \rightarrow 1$ when $(T, \rho) \rightarrow (T_c, \rho_c)$. For the BISM, with small $\mathcal{I}_0$ and δ_J we have $\xi_N^0 \approx a/\pi K_{bcc} \simeq 0.229a$, close to the $d = 3$, nearest neighbor Ising model value [18].

By contrast, the near-critical *charge correlation lengths* depend radically on symmetry. In *charge symmetric models*, where $B_{zz}^z = 1$, it follows from (18) and (24) that the Stillinger-Lovett sum-rule (6) is valid for all fluid regimes *including* the critical point. However, the true charge screening length is given by

$$\xi_{z,\infty}(T, \rho) = R_z(T, \rho) [1 + \mathcal{O}(\mathcal{I}_0^2)], \quad (27)$$

where, from (25) we find $R_z/\xi_D \rightarrow 1$ as $\rho \rightarrow 0$ whereas near criticality one has $R_{z,c}/\xi_{D,c} \approx 2\sqrt{T_0/T_c}$; that yields $R_{z,c}/\xi_{D,c} \simeq 2.36$ for the BISM (for $\mathcal{I}_0$ and δ_J not too large). Furthermore, when $t \rightarrow 0$, the screening length, $\xi_{z,\infty}(\rho_c, T)$ gains, in general, a singular correction factor $[1 + c_{1-\alpha} t^{1-\alpha}]$ [14]. Up to $\mathcal{O}(\mathcal{I}_0)$, the higher moments of S_{zz} satisfy $(\xi_{z,p})^p \approx \xi_D R_z^{p-1}$.

On the other hand, in *nonsymmetric cases* B_{zz}^N in (18) does *not* vanish: rather one has

$$B_{zz}^N(\mathbf{k}; \boldsymbol{\lambda}) = 4\lambda^{\dagger 2} k^4 a^4 / S_d^2 \mathcal{I}_0^2 j_0^2 [1 + \mathcal{O}(k^2)] > 0. \quad (28)$$

Consequently, *all* charge correlations become infected by the divergent density fluctuations controlled by $\Lambda_N(\mathbf{k})$. Nevertheless, because of the factor k^4 , the Stillinger-Lovett relation (6) remains valid in the fluid regime except *at* criticality where it fails and we find [14, 16]

$$(\xi_{z,1}/\xi_D)_c^2 = 1 + w_c^2 \lambda_c^{\dagger 2} = 1 + w_c^2 c_\delta^2 \delta_J^2 + \dots, \quad (29)$$

where $w^2 = 2a^2/S_d \mathcal{I}_0 R_N^2 j_0^2$ and, recalling (23), we note that $c_\delta[J_{\sigma\tau}] \neq 0$ for the BISM. This critical point failure implies a breakdown of full screening that is necessarily associated with slow decay of certain ionic correlations [9]. Indeed, when $(T, \rho) \rightarrow (T_c, \rho_c)$ the charge decay

TABLE I: Charge correlation lengths near criticality where ξ_N diverges, while $R_Z/\xi_D = \mathcal{O}(1)$. The ionicity, $\mathcal{I}_0 \propto q^2$, and asymmetry factor $w_{c\delta}\delta_J$, are defined in (12), (23) and (29).

charge symmetric	non-symmetric
$\xi_{Z,1} = \xi_D = (4\pi\rho q^2/k_B T)^{1/2}$, the Debye length,	$= \xi_D$ for $(T, \rho) \neq (T_c, \rho_c)$, $> \xi_{D,c}$ at (T_c, ρ_c) ,
$\xi_{Z,2}^4 = \xi_D^2 R_Z^2(T, \rho) = \mathcal{O}(\xi_D^4)$,	$= -\xi_D^2 [(w_{c\delta}\delta_J)^2 \xi_N^2 - R_Z^2]$,
$\xi_{Z,p}^{2p} = \xi_D^2 R_Z^{2(p-1)} [1 + \mathcal{O}(\mathcal{I}_0)]$,	$\approx -(w_{c\delta}\delta_J)^2 \xi_D^2 \xi_N^{2(p-1)}$,
$\xi_{Z,\infty} = R_Z [1 + \mathcal{O}(\mathcal{I}_0)] = \mathcal{O}(\xi_D)$,	$\approx \xi_N \sim 1/t^\nu$.

length $\xi_{Z,\infty}$ *asymptotically approaches* the density correlation length, $\xi_{N,\infty}$, and thus *diverges* as ξ_N^0/t^ν (for $\rho = \rho_c$).

However, the fourth charge correlation moment is given (except *at* criticality) by

$$\xi_{Z,2}^4(T, \rho) = \xi_D^2 [R_Z^2 - \xi_{N,1}^2 w^2 \lambda^{\dagger 2}], \quad (30)$$

so that $|\xi_{Z,2}(\rho_c, T)|$ diverges more weakly as $1/t^{\nu/2}$. Note also that on approaching criticality, $\xi_{Z,2}^4$ changes sign (with respect to the symmetric case); for small δ_J this crossover occurs in the BISM at $t = t_\times \approx 0.265 \delta_J$. More generally the higher moments in nonsymmetric systems satisfy $\xi_{Z,p}^{2p} \approx -\xi_{N,\infty}^{2(p-1)} \xi_{D,c}^2 w_c^2 \lambda^{\dagger 2}$ leading to a hierarchy of critical exponents $|\xi_{Z,p}| \sim 1/t^{\nu(1-1/p)}$. Notwithstanding the divergence of the charge correlation length, $\xi_{Z,\infty} \approx \xi_N \rightarrow \infty$, the charge-charge *pair correlation function*, $G_{ZZ}(\mathbf{r})$, decays *exponentially* at (T_c, ρ_c) ! Indeed on approach to criticality we obtain

$$G_{ZZ}(\mathbf{r}) \propto 2\delta_J^2 \frac{T_0}{T_c} \frac{c_\delta^2}{j_0^2} \frac{\xi_D^4}{R_N^2} \frac{e^{-r/\xi_N}}{\xi_N^4(T, \rho)r} - \frac{\xi_D^2}{R_Z^4} \frac{e^{-r/R_Z}}{r}, \quad (31)$$

to leading orders for $d = 3$. At T_c only the second term survives since the first vanishes as $t^{4\nu}$.

Finally, as regards the cross-correlations embodied in the charge-density structure factor S_{NZ} , we find $S_{NZ}(\mathbf{k} = \mathbf{0}; T, \rho) \equiv 0$ except at (T_c, ρ_c) where the value $(\lambda^{\dagger 2} \xi_D^2 / j_0 R_N^2)_c$ is realized. Moreover, on defining moments in analogy to (5), one obtains

$$\xi_{NZ,p}^{2p}(T, \rho) \approx (\lambda^{\dagger} \xi_D^2 / j_0 R_N^2) \xi_{N,\infty}^{2p}(T, \rho), \quad (32)$$

near criticality; of course, all these moments vanish identically in charge symmetric models since $\lambda^{\dagger} \equiv 0$.

In summary, we have analyzed a class of exactly soluble spherical models for 1:1 ionic systems and shown that the Coulomb interactions do not change the gas-liquid critical universality class—contrary to some suggestions [1, 3]. The couplings between the charge correlations and the divergent critical density fluctuations [14, 16] follow mainly from a remarkable structure-function decomposition, Eq. (18), that respects the Stillinger-Lovett (SL)

sum rule (unlike [5]). Our principal results are collected in Table I: they are broadly consistent with Ornstein-Zernike-based arguments advanced for hard-core continuum electrolytes [3]. In *charge-symmetric* models density fluctuations are not directly coupled to two-point charge correlations which remain of short-range and obey SL near and *at* criticality; but in more realistic *nonsymmetric* systems the density fluctuations “infect” the charge correlations which hence exhibit the same diverging correlation length. Moreover, the SL rule is then violated *at* criticality [14, 16] indicating an anomalous conducting state [9].

The authors are grateful to F. Cornu, B. Jancovici, E.R. Smith and G. Stell for correspondence, and to Y. C. Kim for his interest. Support from the NSF (under grants CHE 99-81772 and 03-01101) and assistance to J.-N.A. from the French Ministry of Foreign Affairs under the Lavoisier Fellowship program, is gratefully acknowledged.

-
- [1] See H. Weingärtner and W. Schröer, Adv. Chem. Phys. **116**, 1 (2001) and references therein.
 - [2] M. E. Fisher, J. Stat. Phys. **75**, 1 (1994).
 - [3] G. Stell, J. Stat. Phys. **78**, 197 (1995).
 - [4] B. P. Lee and M. E. Fisher, Phys. Rev. Lett. **76**, 2906 (1996); S. Bekiranov and M. E. Fisher, *ibid* **81**, 5836 (1998).
 - [5] A. Muratov, Sov. Phys.-JETP **93**, 89 (2000).
 - [6] See, e.g., (a) J. M. Romero-Enrique *et al.*, Phys. Rev. Lett. **85**, 4558 (2000); (b) E. Luijten *et al.*, *ibid* **88**, 185701 (2002); (c) Y. C. Kim *et al.*, *ibid* **91**, 065701 (2003).
 - [7] D. C. Brydges and P. Federbush, Commun. Math. Phys. **73**, 197 (1980).
 - [8] F. Stillinger and R. Lovett, J. Chem. Phys. **48**, 3858 (1968).
 - [9] P.-A. Martin, Rev. Mod. Phys. **60**, 1075 (1988).
 - [10] G. S. Joyce, in C. Domb and M.S. Green, Eds, *Phase Transitions and Critical Phenomena*, vol. 2, p.375 (Academic, New York, 1972).
 - [11] (a) E. R. Smith, J. Stat. Phys. **50**, 813 (1988); (b) J. Stat. Phys. [in press].
 - [12] M. C. Barbosa and M. E. Fisher, Phys. Rev. B. **43**, 10635 (1991).
 - [13] F. Cornu and B. Jancovici, J. Stat. Phys. **49**, 33 (1987).
 - [14] J.-N. Aqua and M. E. Fisher : details will be published.
 - [15] A. G. Moreira *et al.*, J. Chem. Phys. **110**, 10058 (1999).
 - [16] Significant differences that arise when $J_{\sigma\tau}^0(\mathbf{R}) \sim 1/R^{d+\varsigma}$, implying $\eta > 0$ when $\varsigma < 2$, will be described elsewhere.
 - [17] Y. C. Kim and M. E. Fisher, J. Phys. Chem. **105**, 11785 (2001).
 - [18] M. E. Fisher and R. J. Burford, Phys. Rev. **156**, 583 (1967).